



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII  
International Rangeland Congress

---

## Comparison of the Acid Detergent Lignin (ADL) among Seven Populations of *lespedeza davurica*

Xiang Zhao  
*Shanxi Agricultural University, China*

Kuanhu Dong  
*Shanxi Agricultural University, China*

Yi Xing  
*Shanxi Agricultural University, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/15-3/2>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Comparison of the acid detergent lignin (ADL) among seven populations of *lespedeza davurica*

Xiang Zhao , Kuanhu Dong , Yi Xing

College of Animal Science and Veterinary Medicine , Shanxi Agricultural University , Taigu , Shanxi , 030801 , China , E-mail : sxndzhaox@126.com

**Key words :** *lespedeza davurica* , population , acid detergent lignin (adl) , cultivation , domestication

**Introduction** *Lespedeza davurica* is an important subshrub in Northern China . It is a drought tolerant , warm season plant . It is grown for hay and pasture , soil improvement , erosion abatement , and to benefit wildlife (Delectis florae reipublicae popularis sinicae agenda academiae sinicae edita , 1995) . Lignification of the plant cell wall has long been considered to be the primary impediment to forage digestibility (Jung , H . G . , 1993) . The purpose of this study was to compare acid detergent lignin (ADL) among seven populations of *Lespedeza davurica* from Henan (HN) and Shanxi (SX) provinces in China .

**Materials and methods** Seven populations (HN and SX) of *Lespedeza davurica* (Table 1) were studied using a randomized complete experimental design , sown using three replicates in a field without irrigation at Jinzhong in Shanxi . Acid detergent lignin (ADL) was measured . (Van Soest , P . J . , and J . B . Robertson . 1980)

**Table 1** The source of the materials .

Code of population	Origin	Longitude latitude
Pop 1	Gong yi city of he nan	112°85' E , 34°26' N
Pop 2	Xin an county of he nan	112°44' E , 34°7' N
Pop 3	Yang quan city of shan xi	113°57' E , 37°85' N
Pop 4	Jiao kou county of shan xi	111°2' E , 36°97' N
Pop 5	Qin shui county of shan xi	114°16' E , 35°70' N
Pop 6	Qin yuan county of shan xi	112°32' E , 36°5' N
Pop 7	Tai gu county of shan xi	112°53' E , 34°72' N

**Results** On the contrary , the content of acid detergent lignin (ADL) increased with procreating (Table 2) . Its value ranged between 5.88% and 8.52% at jointing stage , 10.57% and 13.42% at heading stage , 11.95% and 14.58% at blooming stage , and 12.61% and 17.73% at bearing stage . The values of ADL content were significantly different between the seven populations ( $P < 0.05$ ) . As we already know , at the forepart of growth , the ADL content of grazing was lower and more easily digested because of its tender tress , and at the anaphase , the content was higher and thus harder to digest due to its rough tress .

**Table 2** The acid detergent lignin (ADL) of *Lespedeza davurica* (% of DM) .

Code of population	Jointing stage	Heading stage	Blooming stage	Bearing stage
Pop 1	6.29±0.17de	13.42±1.34a	14.96±0.42a	16.03±0.57b
Pop 2	6.13±0.35e	11.01±0.64bc	11.95±0.54ef	12.61±0.23e
Pop 3	8.52±0.36a	10.57±0.62cd	12.56±0.23cde	13.73±0.71cd
Pop 4	6.76±0.41cd	10.73±0.31bc	12.75±0.76cd	13.66±0.47cd
Pop 5	6.38±0.15de	11.29±0.29bc	12.95±0.67cd	14.31±0.32c
Pop 6	7.60±0.47b	11.86±0.07b	14.58±0.58ab	17.73±0.17a
Pop 7	5.88±0.11e	10.78±0.26bc	13.49±0.14bc	15.80±0.38b

\* In the same column with different small letters means the significantly difference ( $P < 0.05$ )

**Conclusions** Plant cell walls with a high lignin content have long been considered to be the primary impediment to forage digestibility . *Lespedeza davurica* from Xin'an county of Henan was shown to have the lowest ADL , and is used for hay and pasture .

### References

- Delectis florae reipublicae popularis sinicae agenda academiae sinicae edita , 1995 . Flora reipublicae popularis sinicae . Beijing : Science Press . 41 , 131-159 .  
Jung , H . G . , and D . A . Deetz . 1993 . Cell wall lignifications and degradability . 314-315 .